

Patients with confirmed COVID-19 Discontinuing Home Isolation

Patients with confirmed COVID-19 should remain under home isolation precautions until the risk of secondary transmission to others is thought to be low. **The decision to discontinue home isolation precautions should be made on a case-by-case basis, in consultation with healthcare providers and state and local health departments.**

How to discontinue home isolation

- People **with COVID-19 who have stayed home (home isolated)** can stop home isolation under the following conditions:
 - ***If you will not have a test*** to determine if you are still contagious, you can leave home after these three things have happened:
 - You have had no fever for at least 72 hours (that is three full days of no fever without the use medicine that reduces fevers)
AND
 - other symptoms have improved (for example, when your cough or shortness of breath have improved)
AND
 - at least 7 days have passed since your symptoms first appeared
 - ***If you will be tested*** to determine if you are still contagious, you can leave home after these three things have happened:
 - You no longer have a fever (without the use medicine that reduces fevers)
AND
 - other symptoms have improved (for example, when your cough or shortness of breath have improved)
AND
you received two negative tests in a row, 24 hours apart. Your doctor will follow [CDC guidelines](#).

In all cases, **follow the guidance of your healthcare provider and local health department.** The decision to stop home isolation should be made in consultation with your healthcare provider and state and local health departments. Local decisions depend on local circumstances.

Patients with confirmed COVID-19 Discontinuing Home Isolation

For Persons with COVID-19 Under Home Isolation:

The decision to discontinue home isolation should be made in the context of local circumstances. Options now include both 1) a time-since-illness-onset and time-since-recovery (non-test-based) strategy, and 2) a test-based strategy.

Time-since-illness-onset and time-since-recovery strategy (non-test-based strategy)*

Persons with COVID-19 who have symptoms and were directed to care for themselves at home may discontinue home isolation under the following conditions:

- At least 3 days (72 hours) have passed *since recovery* defined as resolution of fever without the use of fever-reducing medications **and** improvement in respiratory symptoms (e.g., cough, shortness of breath); **and**,
- At least 7 days have passed *since symptoms first appeared*.

Test-based strategy (simplified from initial protocol) Previous recommendations for a test-based strategy remain applicable; however, a test-based strategy is contingent on the availability of ample testing supplies and laboratory capacity as well as convenient access to testing. For jurisdictions that choose to use a test-based strategy, the recommended protocol has been simplified so that *only one swab is needed at every sampling*.

Persons who have COVID-19 who have symptoms and were directed to care for themselves at home may discontinue home isolation under the following conditions:

- Resolution of fever without the use of fever-reducing medications **and**
- Improvement in respiratory symptoms (e.g., cough, shortness of breath) **and**
- Negative results of an FDA Emergency Use Authorized molecular assay for COVID-19 from at least two consecutive nasopharyngeal swab specimens collected ≥ 24 hours apart (total of two negative specimens).